

medium-sized anoles may be distinguished in the field by differences in dewlap color and the presence or absence of a white labial stripe (Schwartz and Henderson, 1991). Antillean anoles are often distinguished by color rather than scalation, especially in the hues and tints of the male dewlap (Schwartz, 1969). Differences in scalation between *A. chlorocyanus* and *A. coelestinus* are primarily modal, but differences in the size, shape, and color of dewlap scales are distinctive (Williams, 1965), although some female and small male *A. chlorocyanus* have the pigmented dewlap scales most commonly associated with *A. coelestinus*.

While investigating male agonistic behavior patterns in Barahona, we encountered some individuals that could not be attributed confidently to either species. As a result, we examined 35 green anoles from Barahona and compared them to 22 *Anolis c. chlorocyanus* from Neiba, Provincia de Baoruco (the nearest North Island population from which we had access to an adequate sample), and to 31 *A. c. coelestinus* from the slopes of the Sierra de Baoruco south of Barahona. The two species are clearly allopatric at these two sites. Comparative data are summarized in Table 1. Of the 35 green anoles from Barahona, eight were *A. c. chlorocyanus* and 13 were *A. c. coelestinus*, but 14 had intermediate characteristics indicative of hybrids.

We also observed behavioral evidence indicative of hybridization. On 12 June 1993 near the Hotel Guarocuya in Barahona, we videotaped attempted breeding behavior between a male *Anolis coelestinus* and a female *A. chlorocyanus*. We introduced the tethered female *A. chlorocyanus* to a resident male *A. coelestinus*. Within one minute the male advanced to investigate the female and attempted to mount her. This first encounter lasted approximately 40 sec. The male then retreated up the tree and repositioned himself. We reintroduced the same tethered female to the same male approximately 5 min later. The female responded for about 1 min to the male's advances. The male subsequently succeeded in mounting her, although copulation was not observed. The entire encounter lasted approximately 2.5 min.

Although our data are insufficient to declare with certainty that hybrids exist between these species, 13 of 35 individuals in the zone of contact possessed distinctly intermediate characteristics. This evidence, combined with the lack of hesitation shown in the one interspecific interaction we witnessed, led us to conclude that hybridization occurs. Viability of the purported hybrids apparently is quite high; individuals with intermediate characteristics were among the most aggressive in our studies of male agonistic behavior.

Acknowledgments. -Susan R. Bowersox, Sylvia Calderón, Gretchen L. Cisper, Julio Feliz, Chris Huntington, Lori A. Lenart, Scott P. Sowell, and Kevin C. Zippel helped in the field. Sixto J. and Yvonne A. Incháustegui, Universidad Autónoma de Santo Domingo, José A. Ottenwalder, Parque Zoológico Nacional, and Andreas Schubert, Servicio Alemán, have facilitated opportunities for field work in the Dominican Republic. Permits were provided by Emílio A. Bautista M., Departamento de Viola Silvestre. Field work was supported by Grant No. BIR-9224486 awarded to RP by the National Science Foundation.

LITERATURE CITED

- Powell, R., J. S. Parmerlee, Jr., D. D. Smith, and A. Lathrop. 1991. Geographic distribution: *Anolis chlorocyanus*. *Herpetol. Rev.* 22:134.
- Schwartz, A. 1969. A review of the Hispaniolan lizard *Anolis coelestinus* Cope. *Carib. J. Sci.* 9:33-38.
- and R. W. Henderson. 1991. Amphibians and reptiles of the West Indies: descriptions, distributions, and natural history. Univ. Florida Press, Gainesville, Florida. xvi + 720 pp.
- Williams, E. E. 1965. The species of Hispaniolan green anoles (Sauria: Iguanidae), *Breviora* (227):1-16.

Appendix 1

Specimens examined are in the Bobby Witcher Memorial Collection (BWMC) at Avila College, Kansas City, Missouri 64145 USA: *Anolis c. chlorocyanus* from Neiba, Provincia de Baoruco, República Dominicana (BWMC 02469,03727-9,03734-44, 05378-84); *A. c. coelestinus* from the eastern slopes of the Sierra de Baoruco, Provincia de Barahona, República Dominicana (BWMC 03076-7,03656-73,05268, 05270,05274,05276-7, 05280, 05282-3, 05285-6, 05290); and green anoles from in and near Barahona, Provincia de Barahona, República Dominicana (BWMC 03073-5,04022,04206-11, 04643-4, 04659, 05359-73, 05385-9, 05465, and one specimen being kept alive at Avila College).

Caribbean Journal of Science, Vol. 30, No. 3-4, 281-283, 1994
Copyright 1994 College of Arts and Sciences
University of Puerto Rico, Mayagüez

Records of Bat Ectoparasites from the Luquillo Experimental Forest of Puerto Rico

MICHAEL R. GANNON, *Department of Biology, The Pennsylvania State University, Altoona, Pennsylvania, 16601-3760.*

MICHAEL R. WILLIG, *Ecology Program, Department of Biological Sciences and The Museum, Texas Tech University, Lubbock, Texas 79409-3131.*

During the course of ecological research in the Luquillo Experimental Forest (LEF) of Puerto Rico, we collected ectoparasites from seven species of bats: *Artibeus jamaicensis jamaicensis*, *Brachyphylla cavernarum intermedia*, *Eptesicus fuscus wetmorei*, *Erophylla sezekorni sezekorni*, *Monophyllus redmani portoricensis*, *Pteronotus parnellii portoricensis*, and *Stenoderma rufum darioi*. Herein, we report ectoparasites infesting these bats and provide comments on host-parasite associations.

Bats were captured in mist nets from various locations within the LEF during the dry (March) and rainy (June–August) seasons of 1988 and 1989. Within 20 minutes of capture, wing and tail membranes, pelage, ears, and face were examined for the presence of arthropods. Ectoparasites were removed and placed in vials containing 70% ethyl alcohol; a separate vial stored all ectoparasites from each bat. Hosts were then banded and subsequently released.

TABLE 1. Ectoparasites of bats from Puerto Rico.

Host	Ectoparasite	Family
<i>Artibeus jamaicensis</i>	<i>Megistopoda aranea</i> ^{1,3,7,c}	Streblidae
	<i>Paralabidocarpus foxi</i> ¹	Labidocarpidae
	<i>Paralabidocarpus artibeii</i> ^{4,c}	Labidocarpidae
	<i>Periglischrus iheringi</i> ^{1,2,3,5,7,c}	Spinturnicidae
	<i>Periglischrus vargasi</i> ^{1,2,c}	Spinturnicidae
	<i>Spelaeorhynchus praecursor</i> ^{1,3,6,7,c}	Spelaeorhynchidae
	<i>Aspidoptera phyllostomatus</i> ^{7,a,c}	Streblidae
	<i>Trichobius intermedius</i> ^{1,7,c}	Streblidae
	<i>Trichobius robynae</i> ¹	Streblidae
<i>Brachyphylla cavernarum</i>	<i>Lawrenceocarpus micropilus</i> ^{1,4}	Labidocarpidae
	<i>Lawrenceocarpus puertoricensis</i> ¹	Labidocarpidae
	<i>Radfordiella oudemansi</i> ^{1,3}	Macronyssidae
	<i>Trichobius truncatus</i> ^{1,3,7,c}	Streblidae
	<i>Periglischrus cubanus</i> ^{7,a,b,c}	Spinturnicidae
<i>Eptesicus fuscus</i>	<i>Spinturnix bakeri</i> ^{7,c}	Spinturnicidae
<i>Erophylla sezekorni</i>	<i>Trichobius robynae</i> ¹	Streblidae
	<i>Trichobius truncatus</i> ³	Streblidae
	<i>Ornithodoros viguerasi</i> ³	Argasidae
	<i>Ornithodoros</i> sp. ^{7,c}	Argasidae
	<i>Periglischrus cubanus</i> ^{7,a,c}	Spinturnicidae
<i>Monophyllus redmani</i>	<i>Spelaeorhynchus monophylli</i> ^{1,2,3,6,7,c}	Spelaeorhynchidae
	<i>Trichobius cernyi</i> ^{7,c}	Streblidae
	<i>Trichobius robynae</i> ^{1,7,c}	Streblidae
	<i>Trichobius truncatus</i> ^{1,3}	Streblidae
	<i>Trichobius</i> sp. (near <i>sparsus</i> Kessel) ^{2,c}	Streblidae
	<i>Nycterophilia parnellii</i> ^{7,a,c}	Streblidae
	<i>Periglischrus vargasi</i> ^{7,a,c}	Spinturnicidae
<i>Stenoderma rufum</i>	<i>Paralabidocarpus artibeii</i> ^{1,4,c}	Labidocarpidae
	<i>Paralabidocarpus foxi</i> ^{1,c}	Labidocarpidae
	<i>Paralabidocarpus stenodermi</i> ^{1,c}	Labidocarpidae
	<i>Periglischrus iheringi</i> ^{1,2,3,7,c}	Spinturnicidae
		<i>Cameronieta thomasi</i> ^{7,a,c}

1 = Webb and Loomis, 1977; 2 = Tamsitt and Valdivieso, 1970; 3 = Tamsitt and Fox, 1970a; 4 = Tamsitt and Fox, 1970b; 5 = Rudnick, 1960; 6 = Fain et al., 1967; 7 = this study; a = new record for Puerto Rico; b = new host record; c = known from the Luquillo Experimental Forest,

The primary arthropod groups sampled in this study, spinturnicid wing-mites, spelaeorhynchid ear-mites, argasid ticks, and streblid bat-flies, can be easily seen and collected while examining the bat host. Other ectoparasites may occur on these bats, but were undetected by our sampling protocol,

We collected spinturnicid wing-mites from all the bat species (Table 1), *Periglischrus cubanus* is reported in Puerto Rico for the first time, infesting *Brachyphylla cavernarum* and *Erophylla sezekorni*. This mite was originally described from several species of bats (including *B. nana*) in Cuba (Dusbabek, 1968). *Periglischrus vargasi* infests several species of glossophagine bats, and has been reported from *Monophyllus cubanus* (= *redmani*) in Cuba (Dusbabek, 1968). This is the first report of an association between *P. vargasi* and *M. redmani* in Puerto Rico. *Cameronieta thomasi*, also a

spinturnicid wing-mite, was collected from *Pteronotus parnellii*. *Cameronieta* was first established as a morphologically distinct group from *Periglischrus* by Machado-Allison (1965). This genus appears to be host specific on *Pteronotus* and *Mormoops*, and the association was used to support separation of mormoopid bats from phyllostomids (Smith, 1972). This is the first record of an association between *C. thomasi* and *P. parnellii* in Puerto Rico.

Mites of the family Spelaeorhynchidae were taken from the ears of *Artibeus jamaicensis* (n = 28) and *M. redmani* (n = 9). Unfortunately, all these mites were damaged during removal. The mouthparts of spelaeorhynchids are embedded deeply in the skin, and it is necessary to cut part of the ear to remove them intact. Two species were observed in this study—*Spelaeorhynchus precursor* on *A. jamaicensis*, and *S. mon-*

ophylli on *M. redmani*. These two mites are described and illustrated by Fain et al. (1967).

Argasid ticks (*Ornithodoros* sp.) were collected from *E. sezekorni*. These specimens may be *O. viguerasi*, a taxon first reported on Puerto Rico from *E. bombifrons* (= *sezekorni*) by Tamsitt and Fox (1970a).

Streblid batflies were collected from *A. jamaicensis*, *M. redmani*, and *B. cavernarum*. Of those infesting *A. jamaicensis*, both *Megistopoda aranea* and *Aspidoptera phyllostomatis* are well-known ectoparasites of this host throughout its range. In contrast, associations between *Trichobius intermedius* and *A. jamaicensis* are known only from the Caribbean and Central America (Peterson and Hurka, 1974), including Puerto Rico (Webb and Loomis, 1977). *Trichobius truncatus* was found infesting *B. cavernarum*. This association has also been reported by Tamsitt and Fox (1970a), *Trichobius cernyi* was known previously only from Cuba, and is reported for the first time associated with *M. redmani*. The primary host of *Nycterophilia parnelli* is *Peronotus parnellii*. Although this mormoopid bat is present in Puerto Rico, it was seldom captured during our study. The association of *N. parnelli* on *M. redmani* is a new record for Puerto Rico and may, in this instance, indicate roost sharing between two bat species.

Acknowledgments. —Foremost, we thank R. Waide and A. Lugo for their cooperation and support throughout the duration of this research. We are indebted to A. Estrada-Pinto for providing logistic support. This research was partially supported and performed under grant BSR-8811902 from the National Science Foundation to the Terrestrial Ecology Division, University of Puerto Rico, and the International Institute of Tropical Forestry, as part of the Long-Term Ecological Research Program in the Luquillo Experimental Forest. Additional support was provided by the Forest Service (U.S. Department of Agriculture), the University of Puerto Rico, Sigma Xi, the American Museum of Natural History (Theodore Roosevelt Fund), the Department of Biological Sciences and the Graduate School of Texas Tech University, the American Society of Mammalogists, and Oakridge Associated Universities (U.S. Department of Energy).

LITERATURE CITED

- Dusbabek, F. 1968. Los ácaros cubanos de la Familia Spinturinicidae (Acarina), con notas sobre su especificidad de hospederos. Poeyana, La Habana, Series A, 57:1-31.
- Fain, A., G. Anastos, J. Camin, and D. Johnston. 1967. Notes on the genus *Spelaeorhynchus*, Description of *S. precursor* Neumann and of two new species. *Acarologia* 9:535-556.
- Machado-Allison, C. E. 1965. Notas sobre Mesostigmata Neotropicales. III. *Cameronieta thomasi*: nuevo género y nueva especie parásita de Chiroptera (Acarina, Spinturinicidae). *Acts Biol. Venez.* 4:243-258.
- Peterson, B. V., and K. Hurka. 1974. Ten new species of batflies of the genus *Trichobius* (Diptera: Streblidae). *Can. Entomol.* 106:1049-1066.
- Rudnick, A. 1960. A revision of the mites of the family Spinturinicidae (Acarina), Univ. Calif. Publ. Entomol. 17:157-283.
- Smith, J. D. 1972. Systematic of the chiropteran family Mormoopidae. Misc. Publ. Univ. Kansas Mus. Nat. Hist. 56:1-132.
- Tamsitt, J. R., and I. Fox. 1970a, Records of bat ectoparasites from the Caribbean region (Siphonaptera, Acarina, Diptera). *Can. J. Zool.* 48:1093-1097.
- , and —. 1970b. Mites of the family Lis-trophoridae in Puerto Rico. *Can. J. Zool.* 48:398-399.
- , and D. Valdivieso. 1970. Observations on bats and their ectoparasites. In H. T. Odum and R. F. Pigeon (eds.), *A tropical rain forest*, pp. E123-E128. U.S. Atomic Energy Commission, Oak Ridge, Tennessee.
- Webb, J. P., Jr., and R. B. Loomis. 1977. Ectoparasites. In R. J. Baker et al. (eds.), *Biology of bats of the New World Family Phyllostomatidae, Part II*, pp. 57-119. Spec. Publ., The Museum, Texas Tech Univ. 13:1-364.

Caribbean Journal of Science, Vol. 30, No. 3-4, 283-284, 1994
Copyright 1994 College of Arts and Sciences
University of Puerto Rico, Mayagüez

Biomasa de *Thalassia testudinum* en Punta Gavilán y Santa Cecilia, Quintana Roo, México

ALBERTO DE JESÚS-NAVARRETE, *Centro de Investigaciones de Quintana Roo, A.P. 424 CP 77000, Chetumal Q. Roo, México.*

El pasto marino más estudiado en el mar Caribe es *Thalassia testudinum* (Banks ex König 1805) que ha recibido mucha atención desde el punto de vista biológico (Johnson y Williams, 1982) así como en su ecología y distribución (Fourqurean et al., 1992). En el Caribe Mexicano, la biomasa se ha determinado en la parte norte de Quintana Roo (Nugent et al., 1978; Gallegos et al., 1993), pero no en la zona sur del estado. El objetivo de este trabajo fue determinar la biomasa de *Thalassia testudinum* en dos lugares selectos en la parte sur de Quintana Roo, México.

Se obtuvieron muestras bimensualmente desde febrero a diciembre de 1992 en Punta Gavilán (PG) y Santa Cecilia (SC), al sur de Quintana Roo (18°00'N, 87°45'W y 18°30'N, 88°00'W), Utilizando un cuadrante de 0.1 m se recolectaron muestras al azar a profundidades menores de 1.5 m, dentro de la laguna arrecifal. Todo el material vegetal contenido en el cuadrante fue cortado al nivel del sedimento y se fijó con formol al 4% diluido con agua de mar. En PG se recolectó una muestra y dos réplicas en SC una muestra. Después de eliminar las epífitas y los carbonatos se determinó la biomasa en peso seco (g•ps•m⁻²) (Nugent et al., 1978).